Author Index

Ajima, A.
—, Nakagawa, T. and Kato, T.
Simultaneous measurement of acetylcholine and dopamine releases in rat striatum under freely moving conditions with a brain dialysis method 297
Åkerstedt, T., see Hjemdahl, P. 53
Anderzén, I., see Hjemdahl, P. 53
Ando, S., see Waki, H. 408
Aoki, I., see Miyabayashi, T. 209
Aoyama, T.
—, Kotaki, H. and Saitoh, Y.
Gas chromatographic–mass spectrometric analysis of *threo*-methylphenidate enantiomers in plasma 420
Awni, W.M., see St. Peter, J.V. 424
Beech, J., see Sweeney, R.W. 278
Benjaminsen, T., see Hustvedt, S.O. 335
Benowitz, N.L., see Lee, B.L. 109
Berger, B.J.
—, Hall, J.E. and Tidwell, R.R.
High-performance liquid chromatographic method for the quantification of several diamidine compounds with potential chemotherapeutic value 191
Björkman, S., see Karlsson, A. 157
Bradley, T., see Hjemdahl, P. 53
Canafax, D.M., see Erdmann, G.R. 354
Cappelletti, G., see Guenzi, A. 219
Cassianao, L., see Castagnola, M. 310
Castagnola, M.
—, Cassiano, L., De Cristofaro, R., Luciani, S., Rossetti, D.V. and Landolfi, R.
Purification of the isolated β-chain of adult human haemoglobin from its post-translational modification 310
Castelli, P.L., see Sweeney, R.W. 278
Chan, S.C., see Fox, S.M. 331
Cheng, R., see Kim, C. 67
Chesson, S.M., see Hsu, R.S. 347
Chia, K.S., see Lee, B.L. 303
Chovan, J.P.
— and Vermeulen, J.D.
High-performance liquid chromatographic method for a clozapine analogue, CGS 13429, and its N-oxide and desmethyl metabolites 413
Christophidis, N., see Cosolo, W. 201
Cipolle, R.J., see Erdmann, G.R. 354
Cosolo, W.
—, Drummer, O.H. and Christophidis, N.
Comparison of high-performance liquid chromatography and the Abbott fluorescent polarization radioimmunoassay in the measurement of methotrexate 201
Cristofaro, R. De, see Castagnola, M. 310
Czejka, M.J., see Georgopoulos, A. 340
De Cristofaro, R., see Castagnola, M. 310
Delva, N., see Owen, J.A. 135
Deyl, Z., see Rohlíček, V. 87
DiLeo, E.M., see Hsu, R.S. 347
Drummer, O.H., see Cosolo, W. 201
Dubois, J.P., see Sioufi, A. 361
Dumont, P., see Saraiva Gonçalves, J.C. 389
Ekman, L.
—, Lindström, B. and Roxin, P.
Determination of tacrine and its 1-hydroxy metabolite in plasma using column liquid chromatography with ultraviolet detection 397
Ellis, D.B., see Hsu, R.S. 347
Ellis, P.P., see Wu, P.-Y. 368
Enquist, M.
— and Hermansson, J.
Comparison between two methods for the determination of the total and free (*R*)- and (*S*)-disopyramide in plasma using an α_1-acid glycoprotein column 143
Erdmann, G.R.
—, Gruber, S.A., McGuiggan, M.M., Cripolle, R.J. and Canafax, D.M.
Determination of mizoribine in plasma using ion-pair high-performance liquid chromatography 354
Fischer, G., see Timm, U. 119
Fox, S.M.
—, Chan, S.C. and Van Middlesworth, D.H.
Gas chromatographic and mass spectrometric analysis of articaine in urine 331
Furuta, T., see Kasuya, Y. 101
Gandy, J., see Stidman, J. 318
George, S.R., see Kim, C. 67
Georgopoulos, A.
—, Czejka, M.J., Starzengruber, N., Jäger, W. and Lackner, H.

High-performance liquid chromatographic determination of teicoplanin in plasma: comparison with a microbiological assay 340
Gillberg, M., see Hjemdahl, P. 53
Gleispach, H., see Leis, H.J. 324
Gonçalves, J.C. Saraiva, see Saraiva Gonçalves, J.C. 389
Gruber, S.A., see Erdamnn, G.R. 354
Guenzi, A.
—, Cappelletti, G., Scala, A. and Zanetti, M.
Simultaneous determination of pyrimethamine and mefloquine in human plasma by high-performance liquid chromatography with ultraviolet detection 219
Gupta, K.C., see Kant, K. 289
Hall, J.E., see Berger, B.J. 191
Hall, S.D., see Knedler, M.P. 173
Harvey, D.J.
Long-chain fatty acids and alcohols from gerbil meibomian lipids 23
Hasegawa, R.
—, Murai-Kushiya, M., Komuro, T. and Kumura, T.
Stereoselective determination of plasma pindolol in endotoxin-pretreated rats by high-performance liquid chromatography 381
Heales, S.
— and Hyland, K.
Determination of quinonoid dihydrobiopterin by high-performance liquid chromatography and electrochemical detection 77
Hermansson, J., see Enquist, M. 143
Heuvel, M.P.M. van den, see Onkenhout, W. 31
Hill, H.M., see Selinger, K. 247
Hisaka, A.
—, Kasamatsu, S., Takenaga, N. and Ohtawa, M.
Quantification of L-3-(3-hydroxy-4-pivaloyloxyphenyl)alanine (NB-355) by high-performance liquid chromatography using *o*-phthalaldehyde/N-acetyl-L-cysteine derivatization 183
Hjemdahl, P.
—, Larsson, P.T., Bradley, T., Åkerstedt, T., Anderzén, I., Sigurdsson, K., Gillberg, M. and Lundberg, U.
Catecholamine measurements in urine by high-performance liquid chromatography with amperometric detection - Comparison with an autoanalyser fluorescence method 53
Hsu, R.S.
—, DiLeo, E.M., Chesson, S.M., Ellis, D.B. and Wichmann, J.K.
Determination of velnacrine in plasma and urine by high-performance liquid chromatography 347
Hurtado, M.
—, Medina, M.T., Sotelo, J. and Jung, H.
Sensitive high-performance liquid chromatographic assay for albendazole and its main metabolite albendazole sulphoxide in plasma and cerebrospinal fluid 403
Hustvedt, S.O.
—, Salte, R. and Benjaminsen, T.
Rapid high-performance liquid chromatographic method for the determination of oxolinic acid in fish serum employing solid-phase extraction 335
Hyland, K., see Heales, S. 77
Jacob, III, P., see Lee, B.L. 109
Jäger, W., see Georgopoulos, A. 340
Janicke, D.M., see Jungbluth, G.L. 376
Joshi, A.P., see Kant, K. 289
Jung, H., see Hurtado, M. 403
Jungbluth, G.L.
—, Janicke, D.M. and Jusko, W.J.
Reversed-phase high-performance liquid chromatographic assay for the determination of mezlocillin in human and rat biological samples 376
Jusko, W.J., see Junglbuth, G.L. 376
Kant, K.
-, Joshi, A.P. and Gupta, K.C.
Rapid, quantitative method for the isolation and purification of gangliosides by LIPSEP gel chromatography 289
Karlsson, A.
—, Pettersson, C. and Björkman, S.
Determination of (*R*)- and (*S*)-propranolol in plasma by high-performance liquid chromatography using N-benzoxycarbonylglycyl-L-proline as chiral selector in the mobile phase 157
Kasamatsu, S., see Hisaka, A. 183
Kasuya, Y.
—, Furuta, T. and Shimota, H.
Capillary gas chromatographic–mass

spectrometric determination of stable isotopically labelled and unlabelled theophylline in serum and urine and of 1,3-dimethyluric acid in urine 101
Kataoka, H.
—, Sakiyama, N., Maeda, M. and Makita, M.
Determination of phosphoethanolamine in animal tissues by gas chromatography with flame photometric detection 283
Kato, T., see Ajima, A. 297
Kim, C.
—, Cheng, R. and George, S.R.
Measurement of methionine enkephalin and leucine enkephalin in rat brain regions by high-performance liquid chromatography with coulometric electrochemical detection 67
Kimura, T., see Hasegawa, R. 381
Kitagawa, T., see Matsuura, A. 231
Knadler, M.P.
— and Hall, S.D.
High-performance liquid chromatographic analysis of the enantiomers of flurbiprofen and its metabolites in plasma and urine 173
Komuro, T., see Hasegawa, R. 381
Kotaki, H., see Aoyama, T. 420
Lackner, H., see Georgopoulos, A. 340
Landolfi, R., see Castagnola, M. 310
Larsson, P.T., see Hjemdahl, P. 53
Lawson, J.S., see Owen, J.A. 135
Lee, B.L.
—, Chia, K.S. and Ong, C.N.
Measurement of urinary free catecholamines using high-performance liquid chromatography with electrochemical detection 303
Lee, B.L.
—, Jacob, III, P. and Benowitz, N.L.
Sulfonium salts as derivatizing agents. 2. Determination of theophylline in plasma by automated gas chromatography 109
Leis, H.J.
— and Gleispach, H.
Characterization of the antidiarrhoeal loperamide by gas chromatography–mass spectrometry and application of the Hofmann degradation and Cope elimination reaction 324
Lessard, D., see Selinger, K. 247
Letemendia, F.J.J., see Owen, J.A. 135
Lindström, B., see Ekman, L. 397
Luciani, S., see Castagnola, M. 310
Lundberg, U., see Hjemdahl, P. 53
McGuiggan, M.M., see Erdmann, G.R. 354
Mackert, G.
—, Reinke, M., Schweer, H. and Seyberth, H.W.
Simultaneous determination of the primary prostanoids prostaglandin E_2, prostaglandin $F_{2\alpha}$ and 6-oxoprostaglandin $F_{1\alpha}$ by immunoaffinity chromatography in combination with negative ion chemical ionization gas chromatography–tandem mass spectrometry 13
Maeda, M., see Kataoka, H. 283
Makita, M., see Kataoka, H. 283
Matsuura, A.
—, Nagayama, T. and Kitagawa, T.
Analytical studies on β-lactam antibiotics. III. Automated high-performance liquid chromatographic method for the determination of the orally active antibiotic ceftibuten in human plasma and urine 231
Medina, M.T., see Hurtado, M. 403
Mehta, A.C.
Review. Potential of wide bore open tubular columns in gas chromatographic analysis of drugs 1
Mengel, H., see Nordholm, L. 257
Middlesworth, D.H. Van, see Fox, S.M. 331
Miyabayashi, T.
—, Yamashita, K., Oaki, I., Motohashi, M., Yashiki, T. and Yatani, K.
Determination of manidipine and its pyridine metabolite in human serum by high-performance liquid chromatography with ultraviolet detection and column switching 209
Motohashi, M., see Miyabayashi, T. 209
Murai-Kushiya, M., see Hasegawa, R. 381
Nagayama, T., see Matsuura, A. 231
Nakagawa, T., see Ajima, A. 297
Nordholm, L.
— and Mengel, H.
Determination of an oxadiazole-substituted 1,4-benzodiazepine in plasma by high-performance liquid chromatography with ultraviolet detection and by a radioreceptor assay 257
Ohtawa, M., see Hisaka, A. 183

Ong, C.N., see Lee, B.L. 303
Onkenhout, W.
—, Van der Poel, P.F.H. and Van den Heuvel, M.P.M.
Improved determination of very-long-chain fatty acids in plasma and cultured skin fibroblasts: applications to the diagnosis of peroxisomal disorders 31
Owen, J.A.
—, Sribney, M., Lawson, J.S., Delva, N. and Letemendia, F.J.J.
Capillary gas chromatography of trihexyphenidyl, procyclidine and cycrimine in biological fluids 135
Pappas, A.A., see Stidman, J. 318
Peter, J.V. St., see St. Peter, J.V. 424
Pettersson, C., see Karlsson, A. 157
Poel, P.F.H. van der, see Onkenhout, W. 31
Poupaert, J.H., see Saraiva Gonçalves, J.C. 389
Prelusky, D.B.
—, Warner, R.M. and Trenholm, H.L.
Sensitive analysis of the mycotoxin zearalenone and its metabolites in biological fluids by high-performance liquid chromatography 267
Rassouk, C., see Saraiva Gonçalves, J.C. 389
Reinke, M., see Mackert, G. 13
Riegel, M., see Wu, P.-Y. 368
Rohlíček, V.
— and Deyl, Z.
Simple apparatus for capillary zone electrophoresis and its application to protein analysis 87
Rosseneu, M., see Vercaemst, R. 43
Rossetti, D.V., see Castagnola, M. 310
Roxin, P., see Ekman, L. 397
Saitoh, Y., see Aoyama, T. 420
Sakiyama, N., see Kataoka, H. 283
Salte, R., see Hustvedt, S.O. 335
Sandrenan, N., see Sioufi, A. 361
Saraiva Gonçalves, J.C.
—, Rassouk, C., Poupaert, J.H. and Dumont, P.
High-performance liquid chromatography of chlorambucil prodrugs structurally related to lipids in rat plasma 389
Scala, A., see Guenzi, A. 219
Schweer, H., see Mackert, G. 13
Selinger, K.
—, Lessard, D. and Hill, H.M.
Simultaneous determination of flurazepam and its metabolites in human plasma by high-performance liquid chromatography 247
Seyberth, H.W., see Mackert, G. 13
Shimota, H., see Kasuya, Y. 101
Sigurdsson, K., see Hjemdahl, P. 53
Simmons, H.F., see Stidman, J. 318
Sioufi, A.
—, Sandrenan, N. and Dubois, J.P.
Determination of crotamiton in plasma and urine by high-performance liquid chromatography 361
Sotelo, J., see Hurtado, M. 403
Sribney, M., see Owen, J.A. 135
Starzengruber, N., see Georgopoulos, A. 340
Stidman, J.
—, Taylor, E.H., Simmons, H.F., Gandy, J. and Pappas, A.A.
Determination of meprobamate in serum by alkaline hydrolysis, trimethylsilyl derivatization and detection by gas chromatography–mass spectrometry 318
St. Peter, J.V.
— and Awni, W.M.
Modified high-performance liquid chromatographic assay for antipyrine and its three major metabolites in urine 424
Sweeney, R.W.
—, Beech, J., Whitlock, R.H. and Castelli, P.L.
Analysis of fatty acids in equine cerebrospinal fluid using gas chromatography with electron-capture detection 278
Takenaga, N., see Hisaka, A. 183
Taylor, E.H., see Stidman, J. 318
Tidwell, R.R., see Berger, B.J. 191
Timm, U.
—, Fischer, G., Zell, M. and Zumbrunnen, R.
Determination of the partial benzodiazepine receptor agonist Ro 16-6028 in plasma by capillary gas chromatography with nitrogen-selective detection after conversion into the ethyl ester derivative 119
Trenholm, H.L., see Prelusky, D.B. 267
Union, A., see Vercaemst, R. 43
Van den Heuvel, M.P.M., see Onkenhout, W. 31
Van der Poel, P.F.H., see Onkenhout, W. 31
Van Middlesworth, D.H., see Fox, S.M. 331
Vercaemst, R.
—, Union, A. and Rosseneu, M.

Separation and quantitation of free cholesterol and cholesteryl esters in a macrophage cell line by high-performance liquid chromatography 43
Vermeulen, J.D., see Chovan, J.P. 413
Waki, H.
— and Ando, S.
Column liquid chromatography of calcium channel blockers 408
Warner, R.M., see Prelusky, D.B. 267
Whitlock, R.H., see Sweeney, R.W. 278
Wichmann, J.K., see Hsu, R.S. 347
Wu, P.-Y.
—, Riegel, M. and Ellis, P.P.
High-performance liquid chromatographic assay for timolol in the aqueous humor of the eye 368
Yamashita, K., see Miyabayashi, T. 209
Yashiki, T., see Miyabayashi, T. 209
Yatani, K., see Miyabayashi, T., 209
Zanetti, M., see Guenzi, A. 219
Zell, M., see Timm, U. 119
Zumbrunnen, R., see Timm, U. 119

Subject Index

Acetylcholine
Simultaneous measurement of acetylcholine and dopamine releases in rat striatum under freely moving conditions with a brain dialysis method 297

Albendazole
Sensitive HPLC assay for albendazole and its main metabolite albendazole sulphoxide in plasma and cerebrospinal fluid 403

Albendazole sulphoxide
Sensitive HPLC assay for albendazole and its main metabolite albendazole sulphoxide in plasma and cerebrospinal fluid 403

Alcohols
Long-chain fatty acids and alcohols from gerbil meibomian lipids 23

Antibiotics
Analytical studies on β-lactam antibiotics. III. Automated HPLC method for the determination of the orally active antibiotic ceftibuten in human plasma and urine 231

Antipyrine
Modified HPLC assay for antipyrine and its three major metabolites in urine 424

Articaine
GC and MS analysis of articaine in urine 331

1,4-Benzodiazepine, oxadiazole-substituted
Determination of an oxadiazole-substituted 1,4-benzodiazepine in plasma by HPLC with UV detection and by a radioreceptor assay 257

Benzodiazepine receptor agonist
Determination of the partial benzodiazepine receptor agonist Ro 16-6028 in plasma by capillary GC with nitrogen-selective detection after conversion into the ethyl ester derivative 119

Benzoxycarbonylglycyl-L-proline
Determination of (*R*)- and (*S*)-propranolol in plasma by HPLC using N-benzoxycarbonylglycyl-L-proline as chiral selector in the mobile phase 157

1-[Bis(4-fluorophenyl)methyl]-4-(2,3,4-trimethoxybenzyl)piperazine
Column LC of calcium channel blockers 408

Calcium channel blockers
Column LC of calcium channel blockers 408

Capillary zone electrophoresis
Simple apparatus for capillary zone electrophoresis and its application to protein analysis 87

Catecholamines
Catecholamine measurements in urine by HPLC with amperometric detection – Comparison with an autoanalyser fluorescence method 53

Catecholamines
Measurement of urinary free catecholamines using HPLC with electrochemical detection 303

Ceftibuten
Analytical studies on β-lactam antibiotics. III. Automated HPLC method for the determination of the orally active antibiotic ceftibuten in human plasma and urine 231

CGS 13429
HPLC method for a clozapine analogue, CGS 13429, and its N-oxide and desmethyl metabolites 413

Chlorambucil prodrugs
HPLC of chlorambucil prodrugs structurally related to lipids in rat plasma 389

Cholesterol
Separation and quantitation of free cholesterol and cholesteryl esters in a macrophage cell line by HPLC 43

Cholesteryl esters
Separation and quantitation of free cholesterol and cholesteryl esters in a macrophage cell line by HPLC 43

Cinnarizine
Column LC of calcium channel blockers 408

Clozapine analogue
HPLC method for a clozapine analogue, CGS 13429, and its N-oxide and desmethyl metabolites 413

Columns
Review. Potential of wide bore open tubular columns in GC analysis of drugs 1

Crotamiton
Determination of crotamiton in plasma and urine by HPLC 361
Cycrimine
Capillary GC of trihexyphenidyl, procyclidine and cycrimine in biological fluids 135
N-1-Desalkylflurazepam
Simultaneous determination of flurazepam and its metabolites in human plasma by HPLC 247
Diamidines
HPLC method for the quantification of several diamidine compounds with potential chemotherapeutic value 191
Didesethylflurazepam
Simultaneous determination of flurazepam and its metabolites in human plasma by HPLC 247
3′,4′-Dihydroxyflurbiprofen
HPLC analysis of the enantiomers of flurbiprofen and its metabolites in plasma and urine 173
1,3-Dimethyluric acid
Capillary GC–MS determination of stable isotopically labelled and unlabelled theophylline in serum and urine and of 1,3-dimethyluric acid in urine 101
Disopyramide
Comaparison between two methods for the determination of the total and free (*R*)- and (*S*)-disopyramide in plasma using an α_1-acid glycoprotein column 143
Dopamine
Catecholamine measurements in urine by HPLC with amperometric detection – Comparison with an autoanalyser fluorescence method 53
Dopamine
Simultaneous measurement of acetylcholine and dopamine releases in rat striatum under freely moving conditions with a brain dialysis method 297
Dopamine
Measurement of urinary free catecholamines using HPLC with electrochemical detection 303
Drugs
Review. Potential of wide bore open tubular columns in GC analysis of drugs 1
Enkephalins
Measurement of methionine enkephalin and leucine enkephalin in rat brain regions by HPLC with coulometric electrochemical detection 67
Epinephrine
Cataecholamine measurements in urine by HPLC with amperometric detection – Comparison with an autoanalyser fluorescence method 53
Epinephrine
Measurement of urinary free catecholamines using HPLC with electrochemical detection 303
Fatty acids
Long-chain fatty acids and alcohols from gerbil meibomian lipids 23
Fatty acids
Improved determination of very-long chain fatty acids in plasma and cultured skin fibroblasts: applications to the diagnosis of peroxisomal disorders 31
Fatty acids
Analysis of fatty acids in equine cerebrospinal fluid using GC with electron-capture detection 278
Fatty alcohols
Long-chain fatty acids and alcohols from gerbil meibomian lipids 23
Flunarizine
Column LC of calcium channel blockers 408
Flurazepam
Simultaneous determination of flurazepam and its metabolites in human plasma by HPLC 247
Flurbiprofen
HPLC analysis of the enantiomers of flurbiprofen and its metabolites in plasma and urine 173
Gangliosides
Rapid, quantitative method for the isolation and purification of gangliosides by LIPSEP gel chromatography 289
Haemoglobin, human
Purification of the isolated β-chain of adult human haemoglobin from its post-translational modification 310
4-Hydroxyantipyrine
Modified HPLC assay for antipyrine and

its three major metabolites in urine 424

N-1-Hydroxyethylflurazepam
Simultaneous determination of flurazepam and its metabolites in human plasma by HPLC 247

4′-Hydroxyflurbiprofen
HPLC analysis of the enantiomers of flurbiprofen and its metabolites in plasma and urine 173

3′-Hydroxy-4′-methoxyflurbiprofen
HPLC analysis of the enantiomers of flurbiprofen and its metabolites in plasma and urine 173

Hydroxymethylantipyrine
Modified HPLC assay for antipyrine and its three major metabolites in urine 424

L-3-(3-Hydroxy-4-pivaloyloxyphenyl)alanine
Quantification of L-3-(3-hydroxy-4-pivaloyloxyphenyl)alanine (NB-355) by HPLC using *o*-phthalaldehyde/N-acetyl-L-cysteine derivatization 183

1-Hydroxytacrine
Determination of tacrine and its 1-hydroxy metabolite in plasma using column LC with UV detection 397

β-Lactam antibiotics
Analytical studies on β-lactam antibiotics. III. Automated HPLC method for the determination of the orally active antibiotic ceftibuten in human plasma and urine 231

Leucine enkephalin
Measurement of methionine enkephalin and leucine enkephalin in rat brain regions by HPLC with coulometric electrochemical detection 67

Lipids
Rapid, quantitiative method for the isolation and purification of gangliosides by LIPSEP gel chromatography 289

Long-chain fatty acids
Long-chain fatty acids and alcohols from gerbil meibomian lipids 23

Long-chain fatty alcohols
Long-chain fatty acids and alcohols from gerbil meibomian lipids 23

Loperamide
Characterization of the antidiarrhoeal loperamide by GC-MS and application of the Hofmann degradation and Cope elimination reaction 324

Manidipine
Determination of manidipine and its pyridine metabolite in human serum by HPLC with UV detection and column switching 209

Mefloquine
Simultaneous determination of pyrimethamine and mefloquine in human plasma by HPLC with UV detection 219

Meprobamate
Determination of meprobamate in serum by alkaline hydrolysis, trimethylsilyl derivatization and detection by GC-MS 318

Methionine enkephalin
Measurement of methionine enkephalin and leucine enkephalin in rat brain regions by HPLC with coulometric electrochemical detection 67

Methotrexate
Comparison of HPLC and the Abbott fluorescent polarization radioimmunoassay in the measurement of methotrexate 201

threo-Methylphenidate
GC-MS analysis of *threo*-methylphenidate enantiomers in plasma 420

Mezlocillin
Reversed-phase HPLC assay for the determination of mezlocillin in human and rat biological samples 376

Mizoribine
Determination of mizoribine in plasma using ion-pair HPLC 354

Monodesethylflurazepam
Simultaneous determination of flurazepam and its metabolites in human plasma by HPLC 247

Mycotoxins
Sensitive analysis of the mycotoxin zearalenone and its metabolites in biological fluids by HPLC 267

Norantipyrine
Modified HPLC assay for antipyrine and its three major metabolites in urine 424

Norepinephrine
Catecholamine measurements in urine by HPLC with amperometric detection – Comparison with an autoanalyser fluorescence method 53

Norepinephrine
Measurement of urinary free catechol-

amines using HPLC with electrochemical detection 303

Oxadiazole-substituted 1,4-benzodiazepine
Determination of an oxadiazole-substituted 1,4-benzodiazepine in plasma by HPLC with UV detection and by a radioreceptor assay 257

Oxolinic acid
Rapid HPLC method for the determination of oxolinic acid in fish serum employing solid-phase extraction 335

6-Oxoprostaglandin $F_{1\alpha}$
Simultaneous determination of the primary prostanoids prostaglandin E_2, prostaglandin $F_{2\alpha}$ and 6-oxoprostaglandin $F_{1\alpha}$ by immunoaffinity chromatography in combination with negative ion chemical ionization GC-tandem MS 13

Pentamidine
HPLC method for the quantification of several diamidine compounds with potential chemotherapeutic value 191

Pentamidine analogues
HPLC method for the quantification of several diamidine compounds with potential chemotherapeutic value 191

Peptides
Measurement of methionine enkephalin and leucine enkephalin in rat brain regions by HPLC with coulometric electrochemical detection 67

Phosphoethanolamine
Determanation of phosphoethanolamine in animal tissues by GC with flame photometric detection 283

Pindolol
Stereoselective determination of plasma pindolol in endotoxin-pretreated rats by HPLC 381

Procyclidine
Capillary GC of trihexyphenidyl, procyclidine and cycrimine in biological fluids 135

Propranolol
Determination of (*R*)- and (*S*)-propranolol in plasma by HPLC using N-benzoxycarbonylglycyl-L-proline as chiral selector in the mobile phase 157

Prostaglandins
Simultaneous determination of the primary prostanoids prostaglandin E_2, prostaglandin $F_{2\alpha}$ and 6-oxoprostaglandin $F_{1\alpha}$ by immunoaffinity chromatography in combination with negative ion chemical ionization GC–tandem MS 13

Prostanoids
Simultaneous determination of the primary prostanoids prostaglandin E_2, prostaglandin $F_{2\alpha}$ and 6-oxoprostaglandin $F_{1\alpha}$ by immunoaffinity chromatography in combination with negative ion chemical ionization GC–tandem MS 13

Proteins
Simple apparatus for capillary zone electrophoresis and its application to protein analysis 87

Pyrimethamine
Simultaneous determination of pyrimethamine and mefloquine in human plasma by HPLC with UV detection 219

Quinonoid dihydrobiopterin
Determination of quinonoid dihydrobiopterin by HPLC and electrochemical detection 77

Ro 16-6028
Determination of the partial benzodiazepine receptor agonist Ro 16-6028 in plasma by capillary GC with nitrogen-selective detection after conversion into the ethyl ester derivative 119

Steroids
Separation and quantitiation of free cholesterol and cholesteryl esters in a macrophage cell line by HPLC 43

Tacrine
Determination of tacrine and its 1-hydroxy metabolite in plasma using column LC with UV detection 397

Teicoplanin
HPLC determination of teicoplanin in plasma: comparison with a microbiological assay 340

Theophylline
Capillary GC–MS determination of stable isotopically labelled and unlabelled theophylline in serum and urine and of 1,3-dimethyluric acid in urine 101

Theophylline
Sulfonium salts as derivatizing agents. 2. Determination of theophylline in plasma by automated GC 109

Timolol
HPLC assay for timolol in the aqueous humor of the eye 368
Triethylsulphonium hydroxide
Sulfonium salts as derivatizating agents. 2. Determination of theophylline in plasma by automated GC 109
Trihexyphenidyl
Capillary GC of trihexyphenidyl, procyclidine and cycrimine in biological fluids 135
Velnacrine
Determination of velnacrine in plasma and urine by HPLC 347
Very-long-chain fatty acids
Improved determination of very-long-chain fatty acids in plasma and cultured skin fibroblasts: applications to the diagnosis of peroxisomal disorders 31
Wide bore open tubular columns
Review. Potential of wide bore open tubular columns in GC analysis of drugs 1
Zearalenol
Sensitive analysis of the mycotoxin zearalenone and its metabolites in biological fluids by HPLC 267
Zearalenone
Sensitive analysis of the mycotoxin zearalenone and its metabolites in biological fluids by HPLC 267